## INDEX TO VOLUME II.

ABBOTT, C. C., notice of his 'A Naturalist's Rambles about Home,' 86. Acanthidops bairdi, 291. Accipiter cooperi, 17, 284, fuscus, 17, 174, 284, 311. Actodromas bairdi, 18, 63, 273, fuscicollis, 63, 286. maculata, 18, 286. minutilla, 18, 63, 192, 268. Ægialites, sp., 157. cantianus, 335. melodus, 37. melodus circumcinctus, 285. mongolicus, 240. semipalmatus, 201, 241, 285. vociferus, 375. Ægiothus canescens exilipes, 63. linaria, 16, 99, 200, 240, 279, 291. Æsalon columbarius, 17, 285. richardsoni, 17. 205, 285. Æthyia americana, 10, 18, 288. vallisneria, 18, 288. Agelæus phæniceus, 8, 16, 114, 142, 238, 249, 282. Agersborg, G. S., the birds of Southeastern Dakota, 276. Aix sponsa, 10, 18, 192, 288, 369, 372. Alauda blakistoni, 95. Albatross, Short-tailed, 175. Wandering, 387. Alca cristatella, 341, 342. impennis, 115. psittacula, 342. Alcedo ispida, 95. Aldrich, Charles, a belated bird, Alle nigricans, 211. Allen, J. A, sexual selection and

the nesting of birds, 129; capture

of escaped cage-birds, 314; a Hawk Owl at Chatham, Mass.,

383; a Crested Auk on the Massa-

Aluco flammeus americanus, 284. flammeus pratincola, 192.

chusetts-coast, 388.

American Ornithologists' Union, notes respecting, 223, 317, 318, 389. Ammodramus caudacutus nelsoni, 216, 306. maritimus, 272. savanarum passerinus, 371. Ampelis cedrorum, 15, 279. garrulus, 15, 279. phænicopterum, 291. Amphispiza belli, 16. belli nevadensis, 16. bilineata, 226, 228. Anas americana, 158. boschas, 18, 144, 158, 112, 288, 372. carolinensis, 158. nivalis, 207. obscura, 18, 372. Anorthura fumigata, 291. troglodytes hyemalis, 15, 113. Anser albifrons gambeli, 63, 158, 287hyperboreus nivalis, 207. hutchinsi, 292. leucopareia, 292. Anthus japonicus, 290. ludovicianus, 15, 209, 278, pennsylvanicus, 157, 290, 368. spinoletta, 290, 368. spipoletta, 290, 368. spraguei, 368. Antrostomus carolinensis, 220. Aphelocoma sieberi arizonæ, 371. sordida arizonæ, 356, 371. woodhousei, 16. Aquila chrysaëtus, 205. chrysaëtus canadensis, 10, 17, 157, 285, Archibuteo ferrugineus, 17. lagopus sancti-johannis, 17, 285, 295. Arctonetta fischeri, 158. Ardea cærulea, 272. egretta, 336. herodias, 10, 17, 143, 190, 287.

Arenaria interpres, 157.

Car of the Albert of American Control of the American

Asio accipitrinus, 9, 143, 157, 189, 201, 241, 284.

americanus, 17, 284. wilsonianus, 192.

Astragalinus psaltria, 16, 141. tristis, 16, 141, 279.

Astur atricapillus, 212, 284. atricapillus striatulus, 205. Asturina nitida plagiata, 205. Asyndesmus torquatus, 356.

Atkins, Dr. H. A., notice of death of, 391.

Attila cozumelæ, 295. Auk, Crested, 388.

Great, 115, 209, 374. Paroquet, 388. Razor-billed, 38.

Auriparus flaviceps, 229. Avocet, American, 285, 311. BAGG, Egbert, Jr., the Turkey Buzzard in Central New York, 109.

zard in Central New York, 109. Bain, Francis, the migration of Swallows, 216; a Blue Heron's meal, 221; winter birds of Prince Edward Island, 262.

Baldpate, 288.

Ball, Forrest, Colaptes auratus in

California, 383.
Banks, James W., nest and eggs of the Rusty Grackle (Scolecophagus ferrugineus), 106.

Barrows, W. B., abnormal coloration in a caged Robin, 303; another black Robin, 303.

Bartramia longicauda, 18. 286, 315. Batchelder, Charles F., winter notes from New Mexico, 121, 233; Junco annectens—a correction, 306.

Beckham, Charles W., notes on some of the birds of Pueblo, Colorado, 139; a white-winged Junco in Maryland, 306;

Bell, Robert, notice of his List of Birds observed between Norway House and Forts Churchill and York, 200.

Bernicla brenta, 10, 288.

canadensis, 18, 111, 193, 287. canadensis hutchinsi, 18. canadensis leucopareia, 293. nigricans, 63.

Bicknell, Eugene P., a study of the singing of our birds, 140, 249.

Bird, Myrtle, 99. Bittern, 99, 287, 374.

Least, 101, 296. Blackbird, Brewer's, 282. Crow, 210, 253. Blackbird, Red-and-buff-shoulder'd, 8, 249.

Red-winged, 210, 238, 266, 282.

Rusty, 252, 282.

Yellow-headed, 8, 112, 192, 282, 334.

282, 334. Blackcap, Wilson's, 332. Black-head, Little, 193, 288. Bluebird, 86, 98, 266, 277.

Aretic, 534. California, 330. Mexican, 85, 172.

Rocky Mountain, 8. Bobolink, 97, 98, 152, 189, 266, 282. Bob-white, 45, 315.

Bonasa umbella, 10.

umbella umbelloides, 17, 271. Botaurus lentiginosus, 18, 287. Brachyrhamphus sp., 159. Brant, 10, 288.

Black, 301.

Branta canadensis hutchinsi, 158. Brenta minima, 293.

Brewster, William, Swainson's Warbler, 65; the Heath Hen of Massachusetts, 80; preliminary notes on some birds obtained in Arizona by Mr. F. Stephens in 1884, 84; Swainson's Warbleran omission, 105; Peucæa æstivalis and its subspecies illinoensis, 105; Hawk Owls in New England, 109; the Eider Ducks of the New England coast, 111; the Common Cormorant in Massachusetts, 112; the Rock Ptarmigan of Newfoundland, 193; additional notes on some birds collected in Arizona and the adjoining Province of Sonora, Mexico, by Mr. F. Stephens, with a description of a new species of Ortyx, 196; Nelson's Sharp-tailed Finch (Ammodramus caudacutus nelsoni) on the Atlantic Coast, 216; the Ptarmigan of Anticosti -a correction. 220; additional notes on Swainson's Warbler (Helinaia swainsoni), 346; the

Broad-bill, 193.
Brown, Nathan Clifford, Swamp Sparrows and Yellow-rumps— a question of evidence, 307; the Great Marbled Godwit at Portland, Maine, 385.

liatus) in Massachusetts, 384.

Ovster-catcher (Hamatopus pal-

Browne, F. C., the Red Crossbill breeding in Eastern Massachusetts, 105; the Hawk Owl in Eastern Massachusetts, 220; another Richardson's Owl in Massachusetts, 384.

Bubo virginianus, 212, 284, 298. virginianus arcticus, 17, 21. virginianus saturatus, virginianus subarcticus, 8,

143. Bucephala albeola, 337. islandica, 336. Budytes flavus, 157. Buffle-head, 193. Bunting, Baird's, 268.

Bay-winged, 23. Black-throated, 106, 192, 281,

Henslow's 209. Indigo, 266, 289. Lark, 32, 49, 281. Snow, 101, 265, 279, 290. Varied, 198.

Burgomaster, 37. Buteo borealis, 174, 285. borealis calurus, 9, 17, 356. borealis krideri, 109. borealis socoroensis, 205. brachyurus, 204. cooperi, 17, 165. fuliginosus, 204, 371. lineatus alleni, 292. oxypterus, 371.

swainsoni oxypterus, 371, Bryant, Walter E., the relationship of Podiceps occidentalis and P.

swainsoni, 226, 285.

clarkii, 313. Buzzard, Swainson's, 285. Turkey, 10, 109, 213, 285. CÆLIGENA clemenciæ, 85. 199. Calamospiza bicolor, 16, 32, 226,

281.

melanocorvs, 49. Calcarius Iapponicus, 157 Calipepla californica vallicola, 372. squamata. 213, 226, 228,

Calodromas, 208. Calopezus, 208. Calypte costæ, 231, 309. Campylorhynchus brunneicapillus, 220, 229,

Canace obscura, 10, 17. Cancroma zeledoni, 370. Canvasback, 228. Caprimulgus minor americanus,

178, 179.

Caprimulgus virginianus, 178. vociferus, 257, 272, 283. Cardellina rubrifrons, 353. Cardinal, 72.

Cardinalis cardinalis suberbus, 344. saturatus, 295.

virginianus, 15, 16, 61, 307. Carpodacus cassini, 16, 73, 333, 349. frontalis, 16, 141.

frontalis mexicanus, 207 hæmorrhous, 207. mexicanus, 207. purpureus, 279.

Carpodectes antoniæ, 291. Catbird, 55, 98, 99, 215, 266, 277, 297, 299, 300.

Catharista atrata, 109, 311. Cathartes aura, 10, 17, 109, 143, 285, 356. burrovianus, 293.

urubitinga, 293. Catherpes mexicanus, 233. mexicanus conspersus 15, 350. Cazique, Blue-throated, 85, 199.

Ceophlœus, 52. pileatus, 52.

Centrocercus urophasianus, 10, 17, Centurus carolinus, 17, 283, 335. dubius, 295. uropygialis, 226, 228.

leei, 295. Centronyx bairdi, 16, 198, 231, 333. Centrophanes lapponicus, 16, 63, 190, 280.

ornatus, 16, 280, 333. pictus, 190, 280. Cepphus carbo, 294.

columba, 159, 294. grylle, 294. mandtii, 294. motzfeldi, 294.

Ceratorhina monercerata, 387. Cerorhina suckleyi, 388. Certhia familiaris, 291.

familiaris mexicana, 197, 350. familiaris rufa, 277. familiaris rufa montana, 15. pinus, 343

Certhiola finschi, 293. sancti-thomæ, 293. sundevalli, 293. tricolor, 291.

Ceryle alcyon, 9, 17, 238, 260, 283, superciliosa stictoptera, 208,

Chadbourne, Arthur P., Oporornis agilis and Dendræca palmarum palmarum at Shelburn, New Hampshire, 104; wintering of the Swamp Sparrow in Eastern Massachusetts, 216; Swamp Sparrows and Yellow-rumps, 380.

Chætura pelasgica, 257, 283. yucatanica, 290.

Chamæpelia passerina, 229. Chamberlain, Montague, the nesting habits of the Cape May Warbler (Dendræca tigrina), 33; albino Robins (Turdus migratorius), 102; a white Crow (Corvus frugivorus), 107; the Kingbird in a new rôle, 108; non-appear of Juncos at Montreal, 380

Charadrius dominicus, 18, 285. dominicus fulvus, 157. Chat, Yellow-breasted, 101, 278,

308. Chaulelasmus streperus, 18, 112, 144, 288.

Chen cærulescens, 10, 287. hyperboreus, 18, 287. hyperboreus albatus, 63, 207.

Chewink, 150, 308. Chickadee, Black-capped, 265, 304.

Carolina, 304. Hudsonian, 265. Long-tailed, 277, 330. Mountain, 126, 127, 330.

Chlorostilbon caniveti, 295. forficatus, 295.

Chondestes grammica. 281. grammica strigata, 16, 141.

Chordeiles popetue, 55, 178, 283. popetue henryi, 17, 143. virginianus, 178, 180, 257. virginianus henryi, 180. virginianus minor, 180.

Chroicocephalus franklini, 222. Chrysomitris pinus, 16. 124, 141,

210, 333. Chuck-will's-widow, 211, 220. Cinclus mexicanus, 8, 15, 140, 157,

Circus cyaneus hudsonius, 22, 63,

hudsonius, 9, 17, 157, 284. Cistothorus stellaris, 190, 278.

Clangula albeola, 18, 158, 193, 288. glaucium, 373.

glaucium americana, 18. islandica, 18.

Coale, H. K., a new bird for Illinois, 109.

Coccyzus americanus, 17, 283, 335. erythrophthalmus, 283. Cock, Sage, 10. Colaptes auratus, 259, 284, 335, 383. auratus mexicanus 8, 143. chrysoides, 226, 228, 302. mexicanus, 127, 174, 302, 335, 356.

Colinia, 44.
Colinus, 44, 45.
graysoni, 45.
ridgwayi, 199, 228, 231.
virginianus, 45.
virginianus floridanus, 45.

virginianus texanus, 45. Columba fasciata, 17. 174. Colymbus sp., 158. adamsi, 63.

nigricollis, 340, 341. pacificus, 63. septentrionalis, 63. torquatus, 10, 18, 63, 288.

Contopus albicollis, 290. borealis, 17. depressirostris, 294. pertinax, 356. pileatus, 292. richardsoni, 17, 143. virens, 63, 255, 369.

Conurus carolinensis, 17. Cooke, W. W., notes on the occurrence of certain birds in the Mississippi Valley, 31; Indian bird names, 222.

Coot, American, 10, 112, 287. Cormorant, 210, 211.

Cormorant, Common, 97, 112. Double-crested, 288.

Corvus americanus, 334. corax carnivorus, 9, 16, 157, 174, 282, 371. corax sinuatus, 371. cryptoleucus, 16. frugivorus, 9, 16, 107, 253,

frugivorus, 9, 16, 107, 253. 282.

grebnitzskii, 95.
Cory, C. B., notice of his 'Birds of
Haiti and San Domingo,' 295; notice of his 'List of the Birds of the
West Indies,' 319.

Cotile riparia, 15, 63, 279, 315. Coturniculus henslowi, 190, 209, 281.

lecontei, 23, 190, 381. mexicanus, 370. passerinus, 63, 281, 371. passerinus perpalidus, 16.

Coues. Elliott, probable occurrence of *Diomedea exulans* in Florida, 387.

Cowbird, American, 49, 107, 152, 309.

Cowbird, Bronzed, 211. Crake, Corn, 296, 297. Crane, Sandhill, 10, 287. Whooping, 287. Creeper, Black-and-White, 112, 278.

Brown, 98, 126, 265, 277. Honey, 293.

Mexican, 197 Crex pratensis, 296, 297. Crossbill, American, 8.

Red, 98, 100, 105, 207, 264,

<sup>297</sup>, 333, 379, White-winged, 98, 264. Crow, Clarke's, 101, 282.

Common, 9, 107, 213, 253, 282, 296, 298, 300, 334, 373.

Crymophilus, 182. fulicarius, 182.

Cuckoo, Black-billed, 283. Yellow-billed, 283, 335.

Cupidonia cupido, 17, 82, 285. pinnata, 82.

Curlew, Eskimo, 41, 273, 287. Hudsonian, 10, 287. Long-billed, 10, 41, 273, 286.

Cyanecula suecica, 240. Cyanocitta cristata, 252, 282. stelleri frontalis, 217.

stelleri macrolopha, 16, 125, 142, 174, 355. Cyanocorax cucullatus, 293.

Cyanolesbia, 46. 47. forficata, 47.

mocoa, 47. smaragdina, 47.

Cyclorhis insularis, 295. Cyclorrhynchus psittacula, 159, 388. Cygnus columbianus, 193. Cymbilanius lineatus fasciatus, 294. Cymochorea leucorrhoa, 95, 158. Cypselus saxatilis, 6, 17. Cypseloides niger borealis, 17. Cyrtonyx montezumæ, 46.

DAB-CHICK, 314. Dafila acuta, 18, 63, 193, 201, 241, 288, 372.

Damon, J. L., Cowbird wintering in Western New York, 309.

Davison, J. L., the first nest and eggs of Eremophila alpestris found in Niagara County, N. Y.,

Dendræca æstiva, 15, 141, 278, 331. auduboni, 15, 141, 331, 352. blackburniæ, 103, 376. bryanti. 370. bryanti castaneiceps, 370. cærulea, 15, 278, 369. cærulescens, 62, 65.

coronata, 15, 216, 278. discolor, 62, 278. graciæ, 15, 331, 352. kirtlandi, 376, 381. maculosa, 15, 65, 278, 376. nigrescens, 15, 197, 352. palmarum hypochrysea, 315. palmarum palmarum, 104. petechia rufivertex, 295. pinus, 278, 3.

Dendræca castanea, 65, 272, 315.

rufopileata, 291. striata, 61, 278. tigrina, 33. townsendi, 331. vieilloti, 370. vigorsii, 343.

Dendrocopos immaculatus, 95. Diomedea albatrus, 158. brachyura, 175.

exulans, 387. Dolichonyx oryzivorus, 16, 62, 152, 189. 282.

Dove, Mourning, 10, 285.
Trinidad Ground, 358.
Drew, Frank M., on the vertical

range of birds in Colorado, 11.

Dryobates, 52. Dryopicos, 52. Dryopicus, 52. Dryotomus, 52. Duck, Buffle-head, 337.

Dusky, 336. Eider, 43, 111, 113, 266. Harlequin, 337. Labrador, 209.

Lesser Scaup, 193. Long-tailed, 288, 385. Pin-tail, 193. Redhead, 10, 288. Ring-billed, 288. Ring-neck, 163.

Ruddy, 288. Shoveller, 193, 288. Sprig-tail, 193. Summer, 193.

Wood, 10, 193, 288. Dutcher, William, Bird Notes from Long Island, N. Y., 36; the Can-

ada Goose, 111. Dwight, J., Jr., the Ipswich Spar-row (Passerculus princeps) in

Delaware, 105. Dytes auritus, 96, 289.

Surf, 386.

auritus californicus, 18. nigricollis californicus, 96, 289.

EAGLE, Bald, 10, 192, 285. Golden, 10, 98, 285. White-headed Sea, 192. Ectopistes migratorius, 285. Egret, Great White, 110, 192, 336. Eider, Pacific, 386. Elainea cinerescens, 201. Elanoides forficatus, 19, 21, 192. Elanus leucurus, 205. Empidonax acadicus, 112, 256. atriceps, 291. difficilis, 17. flaviventris, 65, 256. flulvifrons, 207 fulvifrons pallescens, 207. fulvifrons rubicundus, 207. gracilis, 295 hammondi, 17, 143, 283. minimus, 17, 256, 283. obscurus, 17. pusillus, 17. trailli, 256. viridescens, 291. Engyptila gaumeri, 290. Eniconetta stelleri, 158. Ephialtes, 183. Eremophila alpestris, 189, 217. alpestris crysolæma, 17, 239. alpestris leucolæma, 16, 142. Ereunetes occidentalis, 384, 385. pusillus, 18, 63, 143, 201, 241, 286. pusillus occidentalis, 110. Erismatura rubida, 144, 288. Euetheia, 48. bicolor, 48. olivacea intermedia, 395. zena, 48. Eugenes fulgens, 199. FALCO albigularis, 205. arcticus, 188. candicans, 90, 187. candicans β islandicus, 188. candicans gyrfalco, 90, 188. communis, 267. fuscus, 188. grænlandicus, 187, 188. gyrfalco, 90, 187, 188. gyrfalco candicans, 90, 188. gyrfalco grænlandicus, 188. gyrfalco islandicus, 188. holbælli, 187 islandicus, 188, 315. islandicus candicans, 188.

islandus, 185, 187, 188.

islandus β albus, 187. islandus γ maculatus, 187. islandus fuscus, 188.

Falco peregrinus nævius, 9, 17, 212, 285. peregrinus pealei, 157. rusticolus, 185, 188. rusticolus gyrfalco, 187. rusticolus obsoletus, 187. FALCON, Peregrine, 267. Prairie, 285. Finch, Bachman's, 106. Cassin's, 173, 333. Grass, 280. Lark, 281. Lazuli, 281. Nelson's Sharp-tailed, 306. Painted, 72, 298. Pine, 124, 239, 333. Purple, 263, 279. Sea-side, 272, 301. Sharp-tailed, 216, 301, 306. Fisher, A. K., Wood Ibis (*Tantalus loculator*) in Eastern New York, 222; capture of Ammodramus caudacutus nelsoni in the Lower Hudson Valley, New York, 306; capture of two more specimens of Helminthophila leucobronchialis at Sing Sing, New York, 378; evidence concerning the interbreeding of Helminthophila chrysoptera and H. pinus, 378. Flicker, Golden-winged, 284. Flycatcher, Acadian, 112. Arizona Crested, 199. Ash-throated, 98. Black-crested, 242. Great-crested, 254. Hammond's 283. Least, 256, 283. Olivaceus Crested 199. Small Green-crested, 256. Swallow-tailed, 218. Traill's, 256. Wilson's 330 Yellow-bellied, 256. Fratercula corniculata, 159. Frost, Edwin Brant, the Loggerhead Shrike in New Hampshire, Fulica americana, 10, 18 112, 287. caribæa, 292. spinosa, 337 Fringilla æstivalis, 106. bachmani, 105. bicolor, 48, 49. mexicana, 207. zena, 48. Fuligula affinis, 193, collaris, 193.

Fuligula ferina americana, 193, 372. Fulix, affinis, 18, 288. collaris, 18, 158, 288.

marila, 18, 158. Fulmarus glacialis glupischa, 158.

GADWALL, 112, 288. Galeoscoptes carolinensis, 15, 140,

277. Gallinago media wilsoni, 18. wilsoni 286, 336.

Gallinula galeata, 18, 193, 210. Gallinule, Common, 193.

Florida, 193.

Gannet, 98.

Gazetta candidissima, 17, 299. Gault, B. T., nest and eggs of Ca-

lypte costæ, 309. Geococcyx californianus, 17, 143,

Geothlypis bairdi, 294. macgillivrayi, 15, 141, 332.

philadelphia, 65, 278. speciosa, 368. trichas, 15, 62, 65, 278, 367. trichas occidentalis, 141.

Glaucidium phalænoides, 229, 231. Gnat-catcher. Blue-gray, 272, 334.

Godwit, Hudsonian, 286. Marbled, 286, 385.

Golden-eye, 266. Barrows, 336.

Goldfinch, 264, 279. Pine, 264.

oodale, Joseph L., the Wood Thrush in Maine, 215; the Yel-Goodale, low-rumped Warbler wintering in

Maine, 216. Goss, N. S., observations on Elanoides forficatus and Ictinia subcærulea in Kansas, 19; rare summer residents in Kansas, 113; Cyanocitta stelleri frontalis nesting in holes in trees, 217; Wilson's Plover in Nova Scotia, 221; Rissa tridactyla kotzbuei in Washington Territory, 222; the Blackcapped Vireo and Nonpariel in Southwestern Kansas, 274; early accidental occurrence of Catharista atrata and Tantalus loculator in Kansas, 385; the Little Yellow Rail (Porzana novehoraceusis in Kansas, 385; the Thick-billed Grebe (Podylimbus podiceps) Grebe (Podylimbus podiceps) breeding in Kansas, 388. oose, American White-fronted,

Goose,

Canada, 42, 111, 193, 214, 287. Blue-winged, 10, 287.

Goose, Common Wild, 193. Snow, 287

Goshawk, 266, 284 Grackle, Boat-tailed, 100.

Bronzed, 282. Purple, 189.

Rusty, 106, 282, 373. Granatellus sallæi boucardi, 273.

Grebe, American Eared, 96, 289.

Carolina, 96. Clarke's 313. Eared, 340.

European Crested, 93

Horned, 96, 193, 289, 314,

Red-necked 314. Thick-billed, 289, 388.

Western, 313, 314.

Greenlet, 117 Grosbeak, Black-headed, 100, 112, 192, 281.

Blue, 112, 211.

Cardinal, 97, 98, 151, 307,

Evening, 174, 279, 296, 334.

Pine, 264. Rose-breasted, 100, 151, 266,

281, 299. Grouse, Dusky, 10. Pinnated, 80. Ruffed, 10, 40, 270, 300. Sharp-tailed, 285.

Willow, 201. Grus americanus canadensis,

18, 63, 144, 157, 201, 287. Guillemot, Brünnich's, 38. Guiraca cærulea, 16, 112, 142, 211.

Gull, Black-backed, 266. Bonaparte, 266, 374. Burgomaster 37.

Franklin's 289. Glaucous, 37, 337 Herring, 266, 288.

Kittiwake, 266. Pacific Kittiwake, 222. Ring-billed, 288.

Sabine's, 374, 375. Gurney, J. H., notice of his 'List of the Diurnal Birds of Prey,' 201. Gymnocitta cyanocephala, 9, 13, 16.

Gyrfalcon, 187. Gray, 187, 188. Iceland, 284 Labrador 187. White, 187

HÆMATOPUS bachmani, 157. palliatus, 384.

Haliaëtus leucocephalus, 10, 16, 143, 157, 192, 285.

Hardy, Manly, late occurrence of the Phœbe (Sayornis fuscus) at Brewer, Maine, 108. Harelda glacialis, 63, 288. hyemalis, 158, 385.

Harporhynchus cinereus bendirei,

curvirostris palmeri, 196, 228. guttatus, 294.

lecontei, 196, 229, 230, 231. melanostoma, 294. rufus, 15, 140, 277, 298.

Harrier, Common, 22, 192. Hawk, Duck, 9, 209, 285.

Fish, 9, 192, 210, 212, 285. Florida Red-shouldered, 292. Marsh, 9, 100, 192, 285. Pigeon, 100, 285. Red-tailed, 174, 285. Rough-legged, 285, 295. Sharp-skinned, 99, 174, 284, 311.

Sparrow, 100, 285. Western Red-tailed, 9. Hayward, R., curious food for the Kingfisher (Ceryle alcyon), 311.

Helinaia, 76. swainsoni, 76, 105, 346, 377. Helmintherus swainsoni, 301.

vermivorus, 62, 103, 304. Helminthophaga celata, 15, 40, 211,

278, 331. peregrina, 15. ruficapilla, 15. virginiæ, 15, 331.

Helminthophila celata lutescens, 352, 366. chrysoptera, 102, 304, 359,

chrysoptera, 102, 304, 359, 361, 363, 378. gunni, 362. lawrencei, 350, 361, 363.

lawrencei, 359, 361, 363. leucobronchialis, 304, 359, 378, 361. peregrina, 65.

pinus, 359, 361, 363, 378. virginiæ, 352.

Helonæa swainsoni, 62, 104. Hen, Heath, 80, 82.

Prairie, 81, 82, 285.
Henshaw, H. W., the Gulls of the Californian Coast, 231; list of birds observed in summer and fall on the upper Pecos River, 326; a new bird to Massachusetts, 384; the Baird's Sandpiper (Actodromas bairdi) at Locust Grove, New York, 384; the bill of the Horned Puffin (Ceratorhina monocerata), 387.

Herodias alba egretta, 96, 299. egretta, 110, 192.

Heron, Black-crowned Night, 10, 192.
Great Blue, 10, 190, 221, 287.
Green, 287.
Little Blue, 272.
Night, 97, 287.

White, 96, 100, 192. Hesperiphona vespertina, 15, 174,

279, 334, 349. Heteractitis incanus, 157. Hierofalco gyrfalco, 204.

> gyrfalco candicans, 205. gyrfalco islandus, 205, 284. gyrfalco obsoletus, 205. gyrfalco sacer, 63. holbælli, 205.

mexicanus polyagrus, 17, 205, 285.

Himantopus mexicanus, 18. Hirundo erythrogaster, 15, 141, 278. 366.

rustica, 366. rustica erythrogastra, 366. Histrionicus histrionicus, 158, 337, minutus, 18.

Hoffman, W. J., bird names of the Selish, Pah-Uta, and Shoshoni Indians 7

Indians, 7.
Holterhoff, Godfrey, Jr., the Glossy
Ibis and Avocet at San Diego,
Cal., 311

Cal., 311
Hoy, Dr. P. R., nest and eggs of the
Golden-winged Warbler (Helminthopila chrysoptera), 102.

Hummingbird, Black-chinned, 310. Costa, 310. Refulgent, 199. Ruby-throated, 9, 218, 256, 283, 299, 300.

Hydrochelidon lariformis, 211. lariformis surinamensis, 18, 112.

surinamensis, 289.
Hylocichla fuscescens, 277.
fuscescens salicicola, 15.
mustelina, 278, 369.
unalascæ, 330.
unalashæ aububoni, 15, 330.
ustulata swainsoni, 15, 140,

Hylotoma, 51, 52. Hylotomus, 51, 52. pileatus, 283.

IBIS, Glossy, 311. White, 287. Wood, 213, 221, 311.

277, 330.

Icteria virens. 278. virens longicauda, 15, 141. Icterus bullocki, 16, 142, 282. cucullatus, 159, 292. cucullatus igneus, 292. cucullatus nelsoni, 292. curasoënsis, 201. galbula, 16, 55, 58, 251, 282, parisorum, 1, 355 spurius, 16, 251, 282.

Ictinia subcærulea, 19. Indigo-bird, 151.

Ingersoll, Ernest, notice of his 'Country Cousins; Short Studies in the Natural History of the United States,' 91.

Ionornis martinica, 63 Iridoprocne bicolor, 180. JACANA, Brazilian, 337.

Jay, Blue, 100, 101, 253, 265, 282. Blue-fronted, 217. Canada, 99, 107, 265, 335. Long-crested, 125. Piñon, 9.

Steller's, 174.
Jeffries, Wm. A., the Common Cor-morant off Boston Harbor, 112. Jer-Falcon, Brown, 90.

White, 90. Jordan, David Starr, notice of his 'Manual of the Vertebrates of the Northern United States,' 208.

Junco, 115, 116, 117, 380. Mexican, 198 Slate-colored, 146. White-winged, 306.

Junco aikeni, 16, 32. annectens, 16, 306. caniceps, 16, 123, 124, 141. cinereus, 174, 198, 363, 364. cinereus caniceps, 174, 355. cinereus dorsalis, 354, 364. cinereus palliatus, 354, 364 hyemalis, 16, 63, 146, 280, oregonus, 16, 123, 124, 141,

KELLS, Wm. L., foster parents of the Cowbird, 106.

Killdeer, 285. Kingbird, 108, 253, 283. Western, 283.

Kingfisher, 9, 95, 238, 260, 283, 311. Kinglet, Golden-crested, 265, 277. Ruby-crowned, 93, 98, 99, 173,

Kitchen, J. M. W., on the function of the inferior larynx in birds,

Kite, Mississippi, 21.

Kite, Swallow-tailed, 19, 192, 284,

347. Kittiwake, 266. Pacific, 22.

Knot, 312, 375. Lagorus albus, 63, 221. albus alleni, 201. hyperborea, 95. islandorum, 95. lucurus, 14, 17, 95. muta, 94. muta vulgaris, 95. ridgwayi, 95. rupestris, 63, 95, 157, 194, 221.

> ruprestris atkhensis, 95. rupestris nelsoni, 95. rupestris reinhardti, 95, 195. welchi, 194.

Lampornis thalassinus, 295. Lampronetta fischeri, 63, 201. Langille, J. H., notice of his 'Our Birds in their Haunts,' etc., 91. Lantz, D. E., the song of Cardina-

lis virginianus, 307. Lanio melanopygius, 291. Lanius borealis, 15, 212, 279. ludovicianus, 210, 379. ludovicianus excubitorides

15, 279, 332. Lanivireo solitarius plumbeus, 15, 141.

Lark, Horned, 189, 217, 239, 282, Meadow, 250, 266. Mexican Meadow, 198. Shore, 100.

Western Meadow, 8, 282. Larus argentatus smithsonianus, 232, 288.

brachyrhynchus, 232. californicus, 232.

canus, 232. delawarensis, 18, 232, 288. franklini, 288. glaucescens, 158. glaucus, 37, 63. occidentalis, 232. philadelphiæ, 232.

Lawrence, G.N., notice of his paper describing new species of American birds, 290; descriptions of new species of the family Columbidæ,

337. Leucosticte atrata, 16. griseinucha, 157. tephrocotis, 16. tephrocotis littoralis, 16.

Ligea, 295.

Limosa fedoa, 18, 286.
fœda, 385
hæmastica, 286.
lapponica novæ-zelandiæ, 63, 201.
Lobipes hyperboreus, 18, 63, 192, 315.
Lomvia arra, 63.
arra bruennichi, 38.

Long-beak, Greater, 273. Longspur, Chestnut-collared, 280, 333. Lapland, 23, 190, 280, 334.

Lapland, 23, 190, 280, 334. Painted, 190. Smith's, 280.

Loomis, Leverett M., supplementary notes on the ornithology of Chester County, S.C., 168. Loon, 289.

Lohodytes cucullatus, 18, 144, 288. Lohophanes bicolor, 277. inornatus griseus, 15.

wollweberi, 349. Lophortyx californica, 10, 247, 372.

californicus brunnescens, 208, 372.

gambeli, 131, 247, 301. gambeli × californica, 247. Loxia albiventris, 207.

curvirostra americana, 8, 16, 105, 207, 212, 333, curvirostra bendirei, 128, 207, curvirostra mexicana, 207, 372, curvirostra mexicana, 207, 372, curvirostra stricklandi, 372, leucoptera, 16, 210.

pytiopsittacus, 207.
Lunda cirrhata, 159, 388.
M'CALLUM, G. A., albinism, 113.
McKinley, James, field notes from
Pictou County, Nova Scotia, 39.
Macrorhamphus griseus, 18, 374.
griseus scolopaceus, 37, 63,

273, 286. Magpie, Black-billed, 9, 282. Mallard, 100, 112, 289, 336. Mareca americana, 18, 288.

penelope, 373. Martin, Purple, 55, 57, 98, 279, 299,

Megascops, 184. asio, 184. asio bendirei, 184. asio floridanus, 184. asio maccalli, 184. asio maxwelliæ, 184. Megascops flammeolus, 184. trichopsis, 184. Melanerpes erythrocephalus, 17, 63, 143, 259, 283. formicivorus bairdi, 8, 174, 356.

torquatus, 17, 143. Melanetta fusca, 291. Meleagris gallopavo, 10. gallopavo americana, 17, 285. Melopelia leucoptera, 17, 229.

Melospiza cinerea, 157. fasciata, 95, 147, 280. fasciata fallax, 16, 141. fasciata montana, 124, 236, georgiana, 372, lincolni, 16, 65, 142, 280, 354. palustris, 149, 189, 216, 280,

372, 380.
Merganser, 266.
Hooded, 112, 193.
Merganser serrator, 158.
Mergus cucullatus, 193, 372.
merganser, 372.

merganser, 372. mergus americanus, 18, 144, 288. serrator, 201, 241, 372.

Merlin, Richardson's, 285. Merriam, C. Hart, preliminary report of the Committee on Bird Migration, 53; notice of his paper on the Birds of the Bermudas, 95; nest and eggs of the Blackburnian Warbler, 103; Swainson's Warbler off Southern Florida, 104; a remarkable migration of Canada Jays, 107; the White Pelican on Lake Ontario, 111; third addendum to list of birds ascertained to occur within ten miles from Point des Montes, Province of Quebec, Canada; based chiefly upon notes of Napoleon Comeau, 113; Committee on the Migration and Geographical Distribution of North American Birds; Circular for 1885, 117; change of color in the wing-feathers of the Willow Grouse, 201; probable breeding of the Wheatear (Saxicola ananthe) on the northern shore of the Gulf of St. Lawrence, 305; eggs of the Knot (Tringa canutus) found at last, 312; fourth addendum to list of birds ascertained to occur within ten miles of Point de Monts, province of Quebec, Canada; based chiefly upon the notes of Napoleon A. Comeau, 315; Kirtland's Warbler from the Straits of Mackinac, 376; odd nesting habits of the Blue Yellow-backed Warbler in Missouri, 377; Swainson's Warbler in Jamaica, 377; an albino Surf Duck ( Edemia perspicillata), 372.

Merrill, J. C., republication in 'The Auk' of Descriptions of New North American birds,—a suggestion, 389.

Merula merula, 129.

migratoria, 55, 277, 303, 304. migratoria propinqua, 15, 140, 330.

torquata, 129. Micrathene whitneyi, 231.

Microligia, 295. Micropalama himantopus, 273, 286.

Milvulus forficatus, 218. Mimocichla ardesiaca, 205.

Mimus carolinensis, 55, 215, 297. gilvus rostratus, 291.

polyglottus, 15. Minor Ornithological Publications,

96, 209, 295, 372. Mitrephorus pallescens, 207. Mniotilta varia, 15, 62, 278.

Mockingbird, 86, 98. Molothrus æneus, 211.

ater, 16, 49, 50, 106, 142, 282, 309

pecoris, 49, 50, 152.

Motacilla ocularis, 157. Murdoch, John, notes on some species of birds attributed to Point Barrow, Alaska, 200.

Muscicapa atra, 51. carolinensis fusca, 51. fulvifrons, 207.

fusca, 51. nunciola, 51. phæbe, 51.

Myiadestes townsendi, 15, 128, 140, 331.

Myiarchus crinitus, 254. cinerascens, 17, 98.

lawrencei, 208. lawrencei olivaceus, 198, 208. mexicanus magister,

platyrhynchus, 295.

Myiodioctes canadensis, 65. minutus, 372.

pusillus, 278, 330, 332. NAUCLERUS forficatus, 284.

Nelson, E. W., counter-'notes on some species of birds attributed to Point Barrow, Alaska,' 239.

Neocorys spraguei, 269.

Nests and eggs of the birds of Ohio, notice of work on, 289. Nettion carolinensis, 18, 144, 201,

241, 288 Newton, Alfred, notice of his arti-

cle 'Ornithology,' 205. Nighthawk, 55, 178, 283. Cuban, 182.

Western, 182, 283.

Night-jar, 257. Numenius borealis, 63, 273, 287. hudsonicus, 10, 143, 201 241,

> longirostris, 10, 18, 208, 273, 286.

Nutcracker, Clarke's, 282. Nuthatch, Pigmy, 126, 172, 330. Red-bellied, 265, 277.

Slender-billed, 126, 127, 172,

277, 330. White-bellied, 93, 98, 265. Nuttall Ornithological Club, 116,

224, 390. Nutting, Charles C., notice of his paper on a collection of Nicaraguan birds, 293.

Nyctala acadica, 272, 284. tengmalmi richardsoni, 384.

Nyctea nyctea, 157. scandiaca, 37, 63, 284.

Nyctherodius violaceus, 18. Nyctiardea gardeni, 299.

grisea nævia, 10, 192, 287. Nymphicus novæ-hollandiæ, 314. OCEANODROMA furcata, 158. Ochthodromus wilsonius, 221, 273. Œdemia americana, 18, 158.

perspicillata, 158, 386. Œdnicnemus dominicensis, 110. Œstrelata defilippiana, 292.

fisheri, 292. Old-squaw, 266.

Olor americanus, 10. buccinator, 287.

columbianus, 157. Onychotes gruberi, 293.

solitarius, 293. Oreoscoptes montanus, 8, 13, 15,

140 Oriolus ater, 49.

niger, 50. Oriole, Baltimore, 55, 58, 86, 101, 251, 282, 298, 290.

Black, 49. Bullock's 282.

Hooded, 159. Orchard, 251, 282. Scotts, 1.

Ortolan, 190.

Ortygia, 45. Ortygis, 45. Ortyx, 45. Ortyx virginiana, 14, 213, 285, 296, 297, 299, 315. douglasii, 46. elegans, 46. graysoni, 227, 297. massena, 46. montezumæ, 46. picta, 46. Oryzoborus nuttingi, 294. Osgood, Fletcher, the Chuck-will'swidow (Antrostomus carolinensis) in Massachusetts, 220. Osprey, 192. Oxyechus vociferus, 18, 143, 285. Otocoris alpestris, 282. Ouzel, Water, 8, 234, 330. Owl, Acadian, 266. American Long-eared, 9. Barn, 192, 210, 284 Barred, 100, 189, 266, 284. Burrowing, 9, 284. Great Horned, 99, 266, 284. Hawk, 31, 108, 220, 383. Little Screech, 9. Long-eared, 99, 192, 284. Marsh, 189. Monkey-faced, 101. Richardson's, 384. Saw-whet, 272, 284. Screech, 101, 260, 284, 296. Short-eared, 87, 99, 100, 189, Snowy, 37, 366, 284. Western Horned, 9. White, 192. Oyster-catcher, 384. PAGOPHILA eburnea, 63. Palmer, William, abundance of Parus atricapillus near Washington, 304. Pandion haliaëtus, 192. haliaëtus carolinensis, 285. solitarius, 293. Paroquet, Cockatoo, 314. Parra chavaria, 338, dominica, 338, 339. jacana, 339, 338. senegalla, 338, 339. variabilis, 338. Parula americana, 15, 61, 62, 377. nigrilora, 368. pitiayumi, 368. Parus atricapillus, 304. atricapillus septentrionalis,

15, 277.

carolinensis, 304. cinctus, 372, 373. cinctus obtectus, 372. montanus, 15, 126, 127, 140, 331. Passer domesticus, 190, 300. Passerculus alaudinus, 292. anthinus, 212. bairdi, 268. beldingi, 292. princeps, 36, 105. rostratus, 198, 230. sandwichensis, 157. sandwichensis alaudinus, 16, 201, 241. sandwichensis bryanti, 292. sandwichensis savanna, 16, 272, 280. savanna, 65. 376. Passerella iliaca, 150, 281. iliaca schistacea, 16. Passerina amœna, 16, 142, 281. ciris, 276. cyanea, 16, 151, 289. versicolor, 198. Pediœcetes phasianellus campes-tris, 208. phasianellus columbianus, 285. Pelagadroma marina, 386. Pelecanus erythrorhynchus, 18, 111, 144, 288. Pelican, Frigate, 113. White, 111, 288. Pelidna alpina americana, 18, 63. subarquata, 63. Pelionetta perspicillata, 95. Perisoreus canadensis, 335. Perissoglossa tigrina, 32. Petrel, Stormy, 99. White-faced Stormy, 386. Petrochelidon lunifrons, 15, 279, 366. pyrrhonota, 366. Peucæa æstivalis, 106. æstivalis bachmani, 106. æstivalis illinoensis, 105, 106, 299 arizonæ, 226, 370. botteri, 370. carpalis, 228. cassini, 226. mexicana, 370. ruficeps boucardi, 354. Peucedramus olivaceus, 172, 349, 352. Pewee, 51. Wood, 255.

Parus atricapillus turneri, 207, 331.

Phænicothraupis carmioli, 291. Phainopepla nitens, 228, 242. Phalacrocorax carbo, 97, 112, 210. dilophus, 112, 288. dilophus albociliatus, 208. dilophus cincinnatus, 158. pelagicus, 158. Phalænoptilus nuttalli, 9, 17, 143, 283, 356, 382. Phalarope, Northern, 183, 192. Red, 183. Wilson's, 183, 273, 286. Phalaropus fulicarius, 63. lobatus, 157, 183. tricolor, 183. Philacte canagica, 158. Philohela minor, 261, 286. Philomachus pugnax, 336. Philortix, 45. Phloeotomus, 52. Phæbe, 51, 98, 108, 154, 283. Phonipara, 48 Phylloscopus borealis, 373. Pica camtschatica, 95. rustica hudsonica, 9, 16, 142, 282. Picoides arcticus, 210. tridactylis dorsalis, 17. Picicorvus columbianus, 16, 282. Picus insularis, 115. nuttalli, 213. pileatus, 52. pubescens, 257, 283. pubescens gairdneri, 17, 126, 143 stricklandi, 199. villosus, 100, 115, 373. villosus harrisi, 9, 17, 127, 174, 283, 356. Pigeon, Band-tailed, 174. Passenger, 285. Wild. 40. Pinicola enucleator, 15. Pintail, 288. Pittasoma michleri zeledoni, 291. Pipilo aberti, 16. chlorurus, 16, 142. erythrophthalmus, 150, 282, 308, 369. fuscus mesoleucus, 16, 228, 237 maculatus arcticus, 142, 282. maculatus arcticus megalonyx, 16. maculatus megalonyx, 128, Plautus, 183.

Plegadis guarauna, 18.

Plectrophenax hyperboreus, 291. nivalis, 157 Plectrophanes lapponicus, 23, 334. nivalis, 16, 23, 63, 279. ornatus, 23. pictus, 23 Plover, Belted Piping, 285. Black-bellied, 32, 41, 285. Field, 286. Golden, 41, 285. Killdeer, 285, 375. Piping, 37. Semipalmated, 285. Snowy, 335. Upland, 212, 297. Wilson's, 221, 273 Podasocys montanus, 18, 143. Podiceps auritus, 340. californicus, 340. clarkii, 313. cornutus, 193, 314. griseigena holbælli, 314. holbælli, 18. occidentalis, 313, 314. Podilymbus podiceps, 18, 96, 289, 314, 388. Pochard, American, 193. Polioptila cærulea, 15, 272, 334, 349, plumbea, 228. Polysticte stelleri, 63. Poœcetes gramineus, 23, 280, 315. gramineus confinis, 16, 141. Poor-will, 9, 283, 382. Porzana carolina, 18, 62, 190, 209, 287. jamaicensis, 299. leucogastra, 294. noveboracensis, 110, 274, 385. Pratincola rubicola, 129. Procellaria hypoleuca, 387. marina, 387. Progne subis, 15, 55, 57, 228, 279, 299. Protonotaria citrea, 32. Psaltriparus grindæ, 208. minimus californicus, 207. plumbeus, 15. Pseudogryphus californianus, 167. Ptarmigan, Rock, 193, 222. Willow, 221 Puffin, Horned, 387. Pyranga æstiva cooperi, 15, 198, 372. hepatica, 197. ludoviciana, 8, 15, 141, 333, rubra, 95, 278. rubra cooperi, 372.

Pyrrhuloxia sinuata, 228. QUAIL, 285, 296.

Arizona, 296, 299, 301. California, 10, 295. Florida, 45. Grayson's, 45, 297. Masked, 199. Massena, 46. Migratory, 301. Mountain, 300.

Texan, 45 Querquedula carolinensis, 18, 144. 193, 372.

cyanoptera, 112. discors, 18, 144, 288, 372.

Quiscalus, 43. purpureus, 189.

purpureus æneus, 16, 282. RAIL, Carolina, 190.

Clapper, 98, 274. Red-breasted, 287. Sora, 98, 287. Virginia, 96, 98, 192, 287.

Yellow, 274, 385. Rallus elegans, 287. longirostris crepitans 274. virginianus, 18, 144, 192, 287.

Raven, American, 9, 174, 282. Recurvirostra americana, 18, 144, 210, 211, 286.

Redbird, Cardinal, 300. Summer, 98.

Red-head, 193, 288. Redpoll, Lesser, 99, 264, 279. Redstart, American, 98, 278.

Regulus calendula, 14, 15, 173, 277,

japonicus, 291. satrapa, 15, 277, 301.

Ricker, C. B., a specimen of Hel-minthophila leucobronchialis in New Jersey, 378.

Ridgway, Robert, on Buteo har-lani (Aud.) and B. cooperi (Cass.), 165; remarks on the Cal-ifornian Vulture (Pseudogryphus californianus), 167; note on Sarcorhamphus æquatorialis Sharpe, 169; notice of his paper on the American Red Crossbills, 206; notices of his papers on various American birds, 207, 290, 370; do. of his paper on the birds of Cozumel Island, 294; description of a new Cardinal Grosbeak from Arizona, 343; Helmintho-phila leucobronchialis, 359; on Junco cinereus (Swain.) and its geographical races, 363; notice of his paper on the type specimen of Buteo oxypterus, 371; do. emended names of North American birds, 371; a Petrel new to North America, 386; announcement of his nomenclature of Colors and Compendium, etc.

Ridgway Ornithological Club, 116,

Rhodostethia rosea, 63. Rhyacophilus solitarius, 18. Rhynchofalco fusco-cærulescens, 405.

Rhynchophanes maccowni, 16. Rhynchops nigra, 193, 299.

Rissa brevirostris, 158.

tridactyla kotzbuei, 222, 213.

tridactyla pollicaris, 158. Rives, Wm. C., Jr., M. D., nesting of the Worm-eating Warbler (Helmitherus vermivorus) in Virginia, 103; notice of his 'The Birds of Newport,' 208.

Robin 55, 102, 211, 277, 300, 303, 304. Western, 330.

Ruff, 336. SAGE, John H., return of Robins to the same nesting place, 304; another example of Helminthophila leucobronchialis from Connecticut, 304; nesting of the Wormeating Warbler (Helminthotherus vermivorus) in Southern Connecticut, 305; occurrence of the Sharp-shinned Hawk in New Hampshire in winter, 311.

Salpinctes obsoletus, 15, 234, 277. Sandpiper, Baird's, 273, 384.

Bartram's, 375. Bonaparte's, 286. Buff-breasted, 286. Least, 192, 286. Pectoral, 286, 374 Semipalmated, 286. Solitary, 286. Spotted, 286. Stilt. 273, 287. Western Semipalmated, 110,

384, 385. Sarcogrammus cinereus, 339,

goënsis, 339. indicus, 339.

inornatus, 339. Sarcorhamphus æquatorialis, 169.

gryphus, 168, 269, 171. Saunders, W. E., the Black-throated Bunting, Yellow-breasted Chat, and Connecticut Warbler in Ontario, 307.

Saxicola œnanthe, 15, 113, 305. Sayornis, 51.

fusca, 108, 283. phæbe, 51, 254, 369. sayi, 17.

Scardafella inca, 228.

Scolecophagus cyanocephalus, 13, 142, 282.

ferrugineus, 106, 252, 282, 373. Scops, 183.

asio, 9, 184, 260, 284. asio maxwelliæ, 17. flammeolus, 17.

indicus, 184.

Scopus, 184.
Scott, W. E. D., on the breeding habits of some Arizona birds; I, Icterus parisorum, I; II, Icterus cucullatus, 159; III, Phainopepla nitens, 242; IV, Vireo vicinior, 321; winter mountain notes from Southern Arizona, 172; early spring notes from the mountains of Southern Arizona, 348.

of Southern Arizona, 348. Scott, W. L., the Great White Egret and the Yellow Rail in Ottawa, Canada, 110; on the names 'Snowbird' and Winter Wren, 114.

Seebohm, Henry, notice of his 'A History of Britsh Birds,' etc., 88. Se'asphorus flammula, 291.

platycercus, 17. rufus, 17, 356. torridus, 291.

Setophaga picta, 353. ruticilla, 15, 61, 62, 98, 141,

Seton, Ernest E. T., Manitoban Notes, 21; the Swallow-tailed Flycatcher in Manitoba and at York Factory, 218; notes on Manitoban birds, 267; nest and eggs of the Philadelphia Vireo, 305; the Western Grebe in Manitoba, 314; the popular names of birds, 316; interesting records from Toronto, Canada, 334.

Sheldrake, American, 288. Hooded, 288.

Shoveller. 112, 193, 288. Shrike, Great Northern, 238, 278. Loggerhead, 379.

White-rumped, 279, 332.
Shufeldt, R. W., notes of his memoir on the osteology of Ceryle alcyon, 95; do. on the avian patella, 96; variations in the form of the beak, that take place during its growth, in the Short-tailed Al-

batross (Diomedea brachyura), 175; notice of his memoir on the osteology of Numenius longirostris, 208; on the feeding habits of Phalænoptila nuttalli, 382; annoucement of his work entitled 'Contributions to the Anatomy of North American Vertebrates,' 391.

391. Sialia arctica, 8, 15, 140, 234. mexicana, 15, 140, 172, 331,

349. sialis, 15, 140, 277, 381. sialis azurea, 85.

Simorhynchus cristatellus, 159, 341, 388.

psittacula, 159, 342. pygmæus, 159. Sitta canadensis, 15, 277.

aculeatus, 15, 126, 172, 277, 331, 349.

carolinensis, 277, 321, 349. pygmæa, 15, 126, 172, 331, 350.

Siurus auricapillus, 15, 61, 62, 65, 133, 278. motacilla, 91.

nævius, 15, 278. Skimmer, Black, 193. Skylark, Missouri, 369.

Smith, Hugh M., breeding of Loxia americana in the District of Columbia 379; Ereunetes occidentalis on the lower Potomac, 385; Harelda hyemalis in Maryland in summer, 385.

Snipe, American, 336. Red-bellied, 37, 286. Red-breasted, 374. Wilson's 286.

Snowbird, 115, 117.

Black, 98, 265, 280.

White-winged, 32.

Solitaire, Townsend's, 128

Solitaire, Townsend's, 128. Somateria dresseri, 111, 113. mollissima, 111. spectabilis, 63.

v-nigra, 63, 158, 386. Southwick & Jencks, southern range of Rissa tridactyla kotzbuei, 313.

Sparrow, Baird's, 198, 333. Clay-colored, 280. Chipping, 145, 280. English, 190, 211, 296. Field, 145, 280. Fox-colored, 150, 281. House, 99. Harris's, 281. Henslow's, 190. Sparrow, Ipswich, 36, 97, 99, 105.

Java, 314. Large-billed, 198. Leconte's, 23, 190, 280, 381. Lincoln's, 280. Oak-woods, 299. Savanna, 97, 272, 280. Song, 95, 147, 236, 263, 280. Swamp, 100, 149, 189, 216, 280, 307, 380. Titlark, 212.

Tree, 144, 236, 262, 265, 280. White-crowned, 272, 281. White-throated, 281.

Worthen's Field, 291. Yellow-winged, 280. Spatula clypeata, 18, 112, 193, 288. Speotyto cunicularia hypogæa, 9,

17, 143, 284. Spermophila, 47. Sphyropicus thyroides, 17, 174, 349. varius, 17, 258.

varius nuchalis, 9, 17, 174. Spindalis benedicti, 295.

exsul, 295. Spiza americana, 106, 192, 281. Spizella breweri, 16.

domestica, 145, 280. domestica arizonæ, 16, 141. montana, 16. monticola, 144, 280. monticola ochracea, 236. pallida, 16, 141, 280. pusilla, 145, 280. socialis, 380.

wortheni, 291. Sporophila, 47.

morelleti, 47. Squatarola helvetica, 32, 63, 285. Steganopus wilsoni, 18, 273, 286.

Steganopus wilsoni, 18, 273, 286.
Stejneger, Leonard, Analecta ornithologica: XVII, on the origin of the word Quiscalus, 43; XVIII, Colinus, not Ortyx, 44; XIX, is the name Ortyx massena tenable? 46; XX, Cyanolesbia, strictly correct, 46; XXI, concerning the names of some North American Spizinæ, 47; XXII, the correct name of the American Cowbird, 49; XXIII, remarks on the generic name Sayornis and on Sayornis phæbe, 51; XXIV, on the proper generic name of the Pileated Woodpecker and allies, 51; XXV, Why Chordeiles virginianus and not Ch. popetue? 178; XXVI, on the generic names of the Phalaropes, 182; XXVII,

Scops preoccupied, 183; XXVIII, on Gyrfalcons, 184; XXIX, Jacana contra Parra, 337; Colymbus nigricollis, not a North American Bird, 340; XXXI, "Simorhynchus Merrem, 1819," 341; XXXII, Dendroica vigorsii (Aud.), 343; notice of his paper on the Wrens of the subgenus Anorthura, 94; do. the Ptarmigans of the group Attagen, 94; do. descriptions of new species of birds from Kamtschatka. 95; do. list of recent ornithological publications in the United States, 95; notice of his paper on the genus Cepphus, 294; on the alleged occurrence of the Pacific Eider in Labrador, 386.

Stelgidopteryx serripennis, 15, 141,

Stephens, F., notes of an ornithological trip in Arizona and Sonora, 225.

Stercorarius crepidatus, 63. longicaudus, 158. parasiticus, 18, 63, 158, 232. pomatorhinus, 63.

Sterna anæstheta, 230. anglica, 38. antillarum, 289. forsteri, 18, 289. macrura, 63. paradisea, 158.

Strepsilas interpres, 63. Strix nebulosa, 189, 284. Sturnella magna, 250. magna mexicana, 198.

magna mexicana, 198. neglecta, 13, 16, 142, 282. Sula bassana, 98.

Surnia funerea, 31, 108, 220. ulula caparoch, 383. Swallow, Bank, 87, 279.

Barn, 278, 266. Cliff, 279, 366. European Chimney, 366. Rough-winged, 279. Violet-green, 279, 333. White-bellied, 86, 98, 189, 278.

Swan, Trumpter, 287.
Whistling, 10, 193.
Swift, Chimney, 257, 283.
White-throated, 9.

Sylvania microcephala. 372. Sylvia pinus, 343. solitarius, 343.

vigorsii, 343. Symphemia semipalmata, 18, 286, 289. Synthliborhamphus antiquus, 159. TACHYCINETA bicolor, 15, 278. thalassina, 15, 141, 279, 333,

354.
Tachypetes aquila, 113.
Tanager, Cooper's 198.
Liver-colored, 197.
Louisiana, 333.
Scarlet, 278.

Western, 8.
Tantalus loculator, 18, 221, 311.
Teal, Blue-winged, 112, 288, 375.
Green-winged, 193, 238, 288,

Telmatodytes palustris, 15, 190, 278. Tell-tale, Greater, 192.

Lesser, 189. Teretristis fernandina, 368. fornsi, 368.

Tern, Black, 112. Gull-billed, 38.

Thalassidroma pelagica, 99.
Thompson, Frank J., recent occurrence of the Black Vulture in Ohio, 109; the Edicnemus dominicensis in confinement, 110.

Thrasher, Brown, 277. Leconte's, 196. Palmer's, 196. Sage, 8.

Thrush, Brown, 98, 277, 298.
Dwarf, 330.
Golden-crowned, 98, 278.
Long-billed Water, 91.
Olive-backed, 98, 277.
Rocky Mountain Hermit, 330.
Russet-backed, 85.
Small-billed Water, 278.
Swainson's, 330.
Wilson's, 277.

Wood, 72, 215, 277, 288. Thryomanes bewicki leucogaster, 140.

Titlark, 209, 278.

Tinnunculus sparverius, 17, 143, 285.

sparverius isabellinus, 205. Titmouse, Tufted, 277.

Totanus flavipes, 18, 143, 189, 286. melanoleucus, 18, 99, 193, 286. solitarius, 374.

Towhee, 282.

Cañon, 237. Northern, 282. Spurred, 128.

Townsend, Charles H., the occurrence of the Catbird (*Mimus carolinensis*) on the Farallone Islands, Pacific Ocean, 215. Townsend, Charles W., the Blackthroated Bunting in Maine, 106.

Tringa canutus, 63, 312, 375. maculata, 157, 374.

minutilla, 376. Tringoides macularius, 18, 143, 286, 298.

Trochilus alexandri, 310. colubris, 17, 218, 256, 283,

Troglodytes aëdon, 140, 278.
aëdon marianæ, 351.
aëdon parkmani, 15, 278.
alascensis, 94, 157.
beani, 295.
borealis, 94.
fumigatus, 94.
hiemalis, 94.
hiemalis pacificus, 94.
parvulus, 94.
parvulus, 94.

Tryngites rufescens, 63, 286. Turdus migratorius, 102, 114. mustelinus, 215. unalascæ auduboni, 349. ustulatus, 85.

ustulatus aliciæ, 190. Turkey, Wild, 10, 174, 285. Turner, Lucien M., notes on the birds of the Nearer Islands, Alaska, 154; notice of his 'List of the Birds of Labrador,' 368; fa-

miliar Chipping Sparrows, 380. Tyrannula cooperi, 208. mexicana, 208.

Tyrannus carolinensis, 17, 142, 253, 283. tyrannus, 369.

tyrannus, 369. verticalis, 17, 143, 283. vociferans, 17.

URIA grylle, 63. lomvia, 159. Urinator lumme, 159. pacificus, 159. Utamania torda, 38.

Vireo approximans, 291. atricapillus, 274. bairdi, 295. belli. 101. carmioli, 291. gilvus, 279. gilvus swainsoni, 332, 354. huttoni stephensi, 197, 354. magister, 295. noveboracensis, 369.

noveboracensis, 369. olivaceus, 65, 279. philadelphicus, 65, 272, 279, 305.

pusillus, 197.

Vireo solitarius, 65, 279. solitarius cassini, 332. solitarius plumbeus, 332. vicinior, 321.

Vireo, 117. Arizona, 321. Bell's, 101. Blue-headed, 279. Cassin's, 332. Gray, 321. Philadelphia, 272, 279, 305. Plumbeous, 332. Red-eyed, 98, 279. Stephens's, 197. Swainson's, 332. Warbling, 279. White-eyed, 72.

Vireolanius pulchellus verticalis, 293.

cinerea, 295. Vireosylvia cinerea, 295. gilva, 141. grandior, 291.

Vultur californianus, 169. Vulture, Black, 109 California, 167.

WARBLER, Audubon's, 329, 331. Bay-breasted, 272. Black-and-Yellow, 278. Blackburnian, 103. Black-capped, 278. Black-capped Yellow, 278. Black-poll, 278. Black-throated Gray, 197. Black-throated Green, 98. Blue-winged Yellow, 98, 378. Blue Yellow-backed, 97, 377. Cape May, 32, 33. Cerulean, 278. Connecticut, 104, 308. Golden Swamp, 32.

Golden-winged, 102, 378. Grace's, 331. Hooded, 72, 190. Kirtland's, 376. Macgillivray's, 332. Mourning, 278. Olive, 172.

Orange-crowned, 278, 330. Prothonotary, 97.

Prairie, 278. Pine-creeping, 278, 343, 381. Red-poll, 104. Swainson's, 54, 65, 104, 105,

301, 305, 346, 377. Tennessee, 190. Townsend's, 331. White-throated, 378.

Worm-eating, 103, 305.

Warbler, Yellow, 331. Yellow-bellied Red-poll, 190. Yellow-rumped, 98, 216, 278,

307, 380, 381. Yellow-throated, 72.

Waxwing, Cedar, 279. Northern, 279.

Widmann, Otto, note on the capture of Coturniculus lecontei and Dendroæca kirtlandi within the city limits of St. Louis, Mo., 381.

Wilcox, Timothy E., introduced game birds in Oregon and Idaho,

315. Willard, Samuel Wells, food of the Hummingbird (Trochilus colubris), 218; the occurrence of Chroicocephalus franklini in Wisconsin, 222; notice of his paper on the birds of Brown and Outagamie Counties, Wis., 289.

Willet, 286, 298.

Wheatear, 113, 305. Whip-poor-will, 179, 257, 272. Wood, Dr. William, notice of death of, 391.

Woodcock, 99, 213, 261, 286. Woodpecker, Black-backed, 265. Brown-headed, 174.

California, 9, 174. Downy, 101, 257, 265, 283. Gairdner's, 126, 127, Golden-winged, 259, 284, 380.

Hairy, 100, 265. Harris's, 9, 127, 174, 283. Nuchal, 174. Nuttall's, 213.

Pileated, 51, 101, 283. Red-headed, 93, 99, 101, 209. 258, 283.

Red-bellied. 283, 335. Red-naped, 9. Red-shafted, 9, 174. Strickland's, 199. Yellow-bellied, 258, 301.

Wren, Bewick's, 99 Cañon, 233. Carolina. 72, 86, 98, 209. House, 98, 278. Long-billed Marsh, 96, 190,

278. Rock, 234, 277. Short-billed Marsh, 190, 278. Western House, 278.

Winter, 213, 115 XANTHOCEPHALUS icterocephalus, 8, 16, 112, 192, 282.

Xema sabinei, 63. YELLOWBIRD, Summer, 278. Yellow-legs, 286.
Greater, 99, 286.
Yellow-shanks, 189.
Yellow-throat, Maryland, 72, 98, 278, 367.
ZAMELODIA ludoviciana, 151, 192, 281.
melanocephala, 16, 99, 112, 142, 281.
Zeledon, José C., notice of his Catalogue of the Birds of Costa Rica, 370.
Zenaida bogotensis, 358.

Zenaida rubripes, 357.
ruficauda, 291, 357.
vinaceo-rufa, 291, 359.
Zenaidura carolinensis, 9, 17, 143, 285, 372.
macroura, 372.
Zonotrichia albicollis, 65, 281.
gambeli intermedia, 16, 141.
intermedia, 124.
leucophrys, 16, 65, 141, 272, 281.
querula, 281.